



## Confirmation Questions

1. Of those surveyed, what percentage of staff state their Trust has provided specific HIV training?
  - a. 34%
  - b. 23%
  - c. 12%
  - d. 4%
  
2. In the 'friends and family test' survey, a certain percentage of patients state they would not recommend the ambulance service. How much higher is the percentage of HIV+ patients in our survey who state they would not recommend the ambulance service to HIV+ friends and family?
  - a. 19%
  - b. 16%
  - c. 9%
  - d. 3%
  
3. What percentage of people in the UK diagnosed as HIV+ are on treatment?
  - a. 88%
  - b. 96%
  - c. 78%
  - d. 99%
  
4. Life expectancy of people diagnosed with HIV used to be much reduced. Improvements in treatment now mean that their life expectancy has much improved. How many fewer years can someone who is HIV+ expect to live now?
  - a. 5 years
  - b. 1 years
  - c. 1 year
  - d. 0 years
  
5. To be effective, Post-Exposure Prophylaxis (PEP) should be taken within how many hours?
  - a. 24
  - b. 48
  - c. 72
  - d. 120

- 6.** If a patient requires Post-Exposure Prophylaxis (PEP) when sexual health services are not available to provide this, how can they alternatively obtain this?
- a. Prescription via out-of-hours GP
  - b. From pharmacy as an over-the-counter medication
  - c. From A&E
  - d. They cannot, and will need to wait until sexual health service re-opens
- 7.** Beyond universal precautions, what additional precautions should be considered in treating patients who are HIV+?
- a. Double-gloving
  - b. Avoiding invasive procedures except to avoid loss of life
  - c. Ensuring that patient status is handed over as a priority
  - d. None
- 8.** How will someone who is HIV+ know that they are deemed 'undetectable'?
- a. If they are on treatment and feel well
  - b. If they ensure that they take medication regularly as prescribed
  - c. If they have regular viral load monitoring through regular blood tests
  - d. If they have been taking antiretrovirals for at least six weeks
- 9.** In which population group was a reduction in the rate of HIV diagnoses first seen in the UK?
- a. Women
  - b. BME people
  - c. Gay and bisexual men
  - d. IV drug users
- 10.** The ongoing stigma associated with being HIV+ is known to impact on their mental health. What proportion report having suicidal feelings?
- a. 1 in 20
  - b. 1 in 10
  - c. 1 in 5
  - d. 1 in 3